References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

1. Ellis T., Rochester L. Mobilizing Parkinson's Disease: The Future of Exercise. Journal of Parkinson's Disease, vol. 8, no. s1, pp. S95-S100, 2018.

2. Dorsey et al. Projected number of people with Parkinson Disease in the most populous nations, 2005 through 2030. Neurology. 2007: 68(5); 384-6.

3. Huse et al. Burden of illness in Parkinson's Disease. Movement Disorders. 2005: 20(11); 1449-1454.

4. Oguh and Videnovic. Inpatient Management of Parkinson Disease: Current Challenges and Future Directions. The Neurohospitalist. 2012: 2(1); 28-35.

5. Rimmer and Lai. Framing New Pathways in transformative exercise for individuals with existing and newly acquired disability. Disability and Rehabilitation. 2015 Early Online: 1-8.

6. van der Eijk et al. Moving Towards Patient-Centered Healthcare for Patients with Parkinson's Disease. Parkinsonism & Related Disorders. March 2011; 17(5):360-4.

7. Fullard ME, Thibault DP, Hill A, Fox J, Bhatti DE, Burack MA, Dahodwala N, Haberfeld E, Kern DS, Klepitskava OS, Urrea-Mendoza E, Myers P, Nutt J, Rafferty MR, Schwalb JM, Shulman LM, Willis AW, Parkinson Study Group Healthcare Outcomes and Disparities Working Group. 2017; Utilization of rehabilitation therapy services in Parkinson disease in the United States. Neurology 89, 1162-1169.

8. World Health Organization (WHO). International Classification of Functioning, Disability and Health (ICF). https://www.who.int/classifications/icf/en/. Accessed 07/16/19.

9. Keus SHJ, Munneke M, Graziano M, et al. European Physiotherapy Guideline for Parkinson's Disease. 2014; KGNF/ParkinsonNet, the Netherlands.

10. Shin H. and Chung S. Drug-Induced Parkinsonism. Clin Neurol. 2012 Mar; 8(1): 15–21.

11. Parkinson's UK. www.parkinsons.org.uk/information-and-support/exercise-progressing-symptoms.

12. Davis Phinney Foundation. https://b9p3b5u6.stackpathcdn.com/wp-content/uploads/2018/06/Parkinsons-Home-Safety-Checklist-Davis-Phinney-Foundation-2018.pdf. Accessed 08.02.19.

13. Rimmer and Henley. Building the Crossroad between Inpatient/Outpatient Rehabilitation and life-long community based fitness for people with neurologic disability. JNPT 2013;37: 72–77.

14. Rimmer and Lai. Framing New Pathways in transformative exercise for individuals with existing and newly acquired disability. Disability and Rehabilitation. 2015: Early Online: 1–8.

15. Ellis, T., Boudreau, J.K., DeAngelis, T.R. et al. Barriers to exercise in people with Parkinson Disease. PHYS THER: 2013; 93:628-636.

16. Rimmer, J.H., Riley, B., Wang, E. Physical Activity Participation Among persons with Disabilities: Barriers and Facilitators. Am J Prev Med 2004; 26(5): 419-425.

17. Goetz CG, Tilley BC, Shaftman SR, Stebbins GT, Fahn S, Martinez-Martin P, Poewe W, Sampaio C, Stern MB, Dodel R, Dubois B, Holloway R, Jankovic J, Kulisevsky J, Lang AE, Lees A, Leurgans S, LeWitt PA, Nyenhuis D, Olanow CW, Rascol O, Schrag A, Teresi JA, van Hilten JJ, LaPelle N; Movement Disorder

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

Society UPDRS Revision Task Force. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results. Mov Disord. 2008;23(15):2129–70.

18. Espay AJ1, Beaton DE, Morgante F, Gunraj CA, Lang AE, Chen R. Impairments of speed and amplitude of movement in Parkinson's disease: a pilot study. Mov Disord. 2009 May 15;24(7):1001-8. doi: 10.1002/mds.22480.

19. APTA PDEdge recommendations. http://neuropt.org/docs/default-source/parkinson-edge/recommendations-for-patients.pdf?sfvrsn=890d5543_0. Accessed 08.04.19.

20. Gelb DJ, Oliver E, Gilman S. Diagnostic criteria for Parkinson disease. Arch Neurol. 1999 Jan;56(1):33-9.

21. Crispin Jenkinson, Ray Fitzpatrick, Viv Peto, Richard Greenhall & Nigel Hyman (1997) The PDQ-8: Development and validation of a short-form parkinson's disease questionnaire, Psychology & Health, 12:6, 805-814.

22. Schoneburg B, Mancini M, Horak F, Nutt J. Framework for Understanding Balance Dysfunction in Parkinson's Disease. Movement Disorders, Vol. 00, No. 00, 2013.

23. .Duschene C. et al. Influence of aerobic exercise training on the neural correlates of motor learning in Parkinson's disease individuals. Neuroimage Clin. 2016 Sep 14;12:559-569. eCollection 2016.

24.Voss et al. Exercise, brain and cognition across the life span. Journal of Applied Physiology Published 1 November 2011 Vol. 111 no. 5, 1505-1513 DOI: 10.1152/japplphysiol.00210.2011

25.Mang C.S., Snow N.J., Campbell K.L., Ross C.J., Boyd L.A. A single bout of high-intensity aerobic exercise facilitates response to paired associative stimulation and promotes sequence-specific implicit motor learning. J. Appl. Physiol. (1985) 2014;117(11):1325–1336.

26.Dibble LE, Hale TF, Marcus RL, Gerber JP, LaStayo PC. High intensity eccentric resistance training decreases bradykinesia and improves Quality Of Life in persons with Parkinson's disease: a preliminary study. Parkinsonism Relat Disord. 2009 Dec;15(10):752-7. doi: 10.1016/j.parkreldis.2009.04.009. Epub 2009 Jun 3.

27.Dibble LE, Hale T, Marcus RL, Gerber JP, Lastayo PC. The safety and feasibility of high-force eccentric resistance exercise in persons with Parkinson's disease. Arch Phys Med Rehabil. 2006 Sep;87(9):1280-2.

28.Dibble LE, Hale TF, Marcus RL, Droge J, Gerber JP, LaStayo PC. High-intensity resistance training amplifies muscle hypertrophy and functional gains in persons with Parkinson's disease. Mov Disord. 2006 Sep;21(9):1444-52.

29. Bryant M et al. Acute and Long-Term Effects of Multidirectional Treadmill Training on Gait and Balance in Parkinson Disease. PM&R , Volume 8 , Issue 12 , 1151 – 1158.

30. Alberts et al. It Is Not About the Bike, It Is About the Pedaling: Forced Exercise and Parkinson's Disease. Exercise & Sport Sciences Reviews: October 2011 - Volume 39 - Issue 4 - pp 177-186.

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

31. Beall et al. The effect of forced-exercise therapy for Parkinson's disease on motor cortex functional connectivity. Brain Connect. 2013;3(2):190-8. doi: 10.1089/brain.2012.0104. Epub 2013 Feb 25.

32. Ridgel AL1, Vitek JL, Alberts JL. Forced, not voluntary, exercise improves motor function in Parkinson's disease patients. Neurorehabil Neural Repair. 2009 Jul-Aug;23(6):600-8. Epub 2009 Jan 8.

33. Combs et al. Boxing training for patients with Parkinson disease: a case series. Phys Ther. 2011 Jan;91(1):132-42. Epub 2010 Nov 18.

34. Combs et al. Community-based group exercise for persons with Parkinson disease: A randomized controlled trial. NeuroRehabilitation. 2013; vol. 32, no. 1, pp. 117-124.

35. Farley BG, Koshland GF. Training BIG to move faster: the application of the speed-amplitude relation as a rehabilitation strategy for people with Parkinson's disease. Exp Brain Res. 2005 Dec;167(3):462-7. Epub 2005 Nov 11.

36. Ebersbach G et al. Comparing exercise in Parkinson's disease--the Berlin LSVT®BIG study. Mov Disord. 2010 Sep 15;25(12):1902-8. doi: 10.1002/mds.23212. Erratum in: Mov Disord. 2010 Oct 30;25(14):2478.

37. Fox C, Ebersbach G, Ramig L, Sapir S. LSVT LOUD and LSVT BIG: Behavioral Treatment Programs for Speech and Body Movement in Parkinson Disease. Parkinsons Dis. 2012;2012:391946. Epub 2012 Mar 15.

38. Janssens J, Malfroid K, Nyffeler T, Bohlhalter S, Vanbellingen T. Application of LSVT BIG intervention to address gait, balance, bed mobility, and dexterity in people with Parkinson disease: a case series. Phys Ther. 2014 Jul;94(7):1014-23. doi: 10.2522/ptj.20130232. Epub 2014 Feb 20.

39. Ebersbach G, Grust U, Ebersbach A, Wegner B, Gandor F, Kühn AA. Amplitude-oriented exercise in Parkinson's disease: a randomized study comparing LSVT-BIG and a short training protocol. J Neural Transm (Vienna). 2015 Feb;122(2):253-6. Epub 2014 May 29.

40. APTA PDEdge recommendations. http://neuropt.org/docs/default-source/parkinson-edge/recommendations-for-patients.pdf?sfvrsn=890d5543_0. Accessed 08.04.19.

41. Gelb DJ, Oliver E, Gilman S. Diagnostic criteria for Parkinson disease. Arch Neurol. 1999 Jan;56(1):33-9.

42. Colgrove et al. Effect of Yoga on Motor Function in People with Parkinson's Disease: A Randomized, Controlled Pilot Study. Colgrove et al., J Yoga Phys Ther 2012, 2:2.

43. Roland, K. Applications of yoga in Parkinson's disease: a systematic literature review. Journal of Parkinsonism and Restless Legs Syndrome. 9 January 2014

44. Li F, Harmer P, Fitzgerald K, et al. Tai chi and postural stability in patients with Parkinson's disease. N Engl J Med. 2012;366: 511–519.

45. Hackney ME, Earhart GM. Tai Chi improves balance and mobility in people with Parkinson disease. Gait Posture 2008;28(3):456–60.

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

46. Keus SH, Bloem BR, Hendriks EJ, et al. Evidence-based analysis of physical therapy in Parkinson's disease with recommendations for practice and research. Mov Disord. 2007;22:451–460.

47. Schenkman, M, Cutson, TM, Kuchibhatla, M, et al. Exercise to improve spinal flexibility and function for people with Parkinson's disease: a randomized, controlled trial. J Am Geriatr Soc. 1998;46:1207–1216.

48. Hughes AJ, Daniel SE, Blankson S, Lees AJ. A clinicopathologic study of 100 cases of Parkinson disease. Arch Neurol. 1993;50140- 148.

49. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. J Neurol Neurosurg Psychiatry. 1992;55181- 184

50. Rajput AH, Rozdilsky B, Rajput A. Accuracy of clinical diagnosis in parkinsonism: a prospective study. Can J Neurol Sci. 1991;18275- 278.

51. APTA PDEdge recommendations. http://neuropt.org/docs/default-source/parkinson-edge/recommendations-for-patients.pdf?sfvrsn=890d5543_0. Accessed 08.04.19.

52. Tremor management for Parkinson's Disease and Essential Tremors. UF center for movement disorders and neurorestoration. http://movementdisorders.ufhealth.org/2015/02/12/tremor-management-for-parkinsons-disease-and-essential-tremors/. Accessed June 22, 2017.

53. Westheimer O, McRae C, Henchcliffe C, Fesharaki A, Glazman S, Ene H, Bodis-Wollner I. J Neural Transm (Vienna). 2015 Sep;122(9):1263-70.

54. McGill A, Houston S, Lee RY. Dance for Parkinson's: a new framework for research on its physical, mental, emotional, and social benefits. Complement Ther Med. 2014 Jun;22(3):426-32.

55. Alberts J et al. It Is Not About the Bike, It Is About the Pedaling: Forced Exercise and Parkinson's Disease. Exercise & Sport Sciences Reviews: October 2011 - Volume 39 - Issue 4 - pp 177-186.

56. Beall et al. The effect of forced-exercise therapy for Parkinson's disease on motor cortex functional connectivity. Brain Connect. 2013;3(2):190-8. doi: 10.1089/brain.2012.0104. Epub 2013 Feb 25.

57. Ridgel AL1, Vitek JL, Alberts JL. Forced, not voluntary, exercise improves motor function in Parkinson's disease patients. Neurorehabil Neural Repair. 2009 Jul-Aug;23(6):600-8. Epub 2009 Jan 8.

58. Allen NE, Sherrington C, Paul SS, Canning CG. Balance and falls in Parkinson's disease: a meta-analysis of the effect of exercise and motor training. Mov Disord. 2011;26:1605–1615.

59. Schoneburg B, Mancini M, Horak F, Nutt JG. Framework for understanding balance dysfunction in Parkinson's disease. Mov Disord 2013; 28: 1474–1482.

60. Boonstra TA, van Kordelaar J, Engelhart D, van Vugt JP, van der Kooij H. Asymmetries in reactive and anticipatory balance control are of similar magnitude in Parkinson's disease patients. Gait Posture 2016; 43: 108–113.

61. Smulders K, Esselink RA, De Swart BJ, Geurts AC, Bloem BR, Weerdesteyn V. Postural inflexibility in PD: does it affect compensatory stepping? Gait Posture 2014; 39: 700–706.

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

62. Duncan RP, Leddy AL, Cavanaugh JT, Dibble LE, Ellis TD, Ford MP, et al. Balance differences in people with Parkinson disease with and without freezing of gait. Gait Posture 2015; 42: 306–309. 11.

63. Vervoort G, Bengevoord A, Strouwen C, Bekkers EM, Heremans E, Vandenberghe W, et al. Progression of postural control and gait deficits in Parkinson's disease and freezing of gait: a longitudinal study. Parkinsonism Relat Disord 2016; 28: 73–79.

64. Bloem BR, Marinus J, Almeida Q, Dibble L, Nieuwboer A, Post B, Ruzicka E, Goetz C, Stebbins G, Martinez-Martin P, Schrag A; Movement Disorders Society Rating Scales Committee. Measurement instruments to assess posture, gait, and balance in Parkinson's disease: Critique and recommendations. Mov Disord. 2016 Sep;31(9):1342-55. doi: 10.1002/mds.26572. Epub 2016 Mar

65. Schoneburg B, Mancini M, Horak F, Nutt J. Framework for Understanding Balance Dysfunction in Parkinson's Disease. Movement Disorders, Vol. 00, No. 00, 2013.

66. Bloem BR, Beckley DJ, van Dijk JG. Are automatic postural responses in patients with Parkinson's disease abnormal due to their stooped posture? Exp Brain Res. 1999;124:481–488.

67. Jacobs JV, Dimitrova DM, Nutt JG, et al. Can stooped posture explain multidirectional postural instability in patients with Parkinson's disease? Exp Brain Res. 2005;166:78–88.

68. Forsyth AL, Paul SS, Allen NE, Sherrington C, Fung VS, Canning CG. Flexed Truncal Posture in Parkinson Disease: Measurement Reliability and Relationship With Physical and Cognitive Impairments, Mobility, and Balance. J Neurol Phys Ther. 2017 Apr;41(2):107-113.

69. Mancini M, Horak FB, Zampieri C, et al. Trunk accelerometry reveals postural instability in untreated Parkinson's disease. Parkinsonism Relat Disord. 2011;17:557–562.

70. Stylianou AP, McVey MA, Lyons KE, et al. Postural sway in patients with mild to moderate Parkinson's disease. Int J Neurosci. 2011;121:614–621.

71. Nantel J, McDonald JC, Bronte-Stewart H. Effect of medication and STN-DBS on postural control in subjects with Parkinson's disease. Parkinsonism Relat Disord. 2012;18:285–289.

72. Mitchell SL, Collins JJ, De Luca CJ, et al. Open-loop and closedloop postural control mechanisms in Parkinson's disease: increased mediolateral activity during quiet standing. Neurosci Lett. 1995; 197:133–136. 22.

73. Matinolli M, Korpelainen JT, Korpelainen R, et al. Postural sway and falls in Parkinson's disease: a regression approach. Mov Disord. 2007;22:1927–1935.

74. Frenklach A, Louie S, Koop MM, et al. Excessive postural sway and the risk of falls at different stages of Parkinson's disease. Mov Disord. 2009;24:377–385.

75. Rocchi L, Chiari L, Horak FB. Effects of deep brain stimulation and levodopa on postural sway in Parkinson's disease. J Neurol Neurosurg Psychiatry. 2002;73:267–274.

76. Bronte-Stewart HM, Minn AY, Rodrigues K, et al. Postural instability in idiopathic Parkinson's disease: the role of medication and unilateral pallidotomy. Brain. 2002;125:2100–2114. 17. Colnat-Coulbois S,

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

Gauchard GC, Maillard L, et al. Bilateral subthalamic nucleus stimulation improves balance control in Parkinson's disease. J Neurol Neurosurg Psychiatry. 2005;76:780– 787.

77. Colnat-Coulbois S, Gauchard GC, Maillard L, et al. Bilateral subthalamic nucleus stimulation improves balance control in Parkinson's disease. J Neurol Neurosurg Psychiatry. 2005;76:780– 787.

78. Peterka RJ, Benolken MS. Role of somatosensory and vestibular cues in attenuating visually-induced human postural sway. In: Woollacott M, Horak F, eds. Posture and Gait: Control Mechanisms. Eugene, OR: University of Oregon Press; 1992:272–275.

79. Horak FB, Frank J, Nutt J. Effects of dopamine on postural control in parkinsonian subjects: scaling, set, and tone. J Neurophysiol. 1996;75:2380–2396.

80. Jacobs JV, Tran V, Nutt JG, et al. Push and release: an alternative clinical test of postural instability for patients with Parkinson's disease. J Neurol. 2006;77:322–326.

81. King LA, St George RJ, Carlson-Kuhta P, et al. Preparation for compensatory forward stepping in Parkinson's disease. Arch Phys Med Rehabil. 2010;91:1332–1338.

82. Hallett M, Khoshbin S. A physiological mechanism of bradykinesia. Brain. 1980;103:301–314.

83. Bloem BR, Marinus J, Almeida Q, Dibble L, Nieuwboer A, Post B, Ruzicka E, Goetz C, Stebbins G, Martinez-Martin P, Schrag A; Movement Disorders Society Rating Scales Committee. Measurement instruments to assess posture, gait, and balance in Parkinson's disease: Critique and recommendations. Mov Disord. 2016 Sep;31(9):1342-55. doi: 10.1002/mds.26572. Epub 2016 Mar 4.

84. Allen NE, Sherrington C, Paul SS, Canning CG. Balance and falls in Parkinson's disease: a meta-analysis of the effect of exercise and motor training. Mov Disord. 2011;26:1605–1615.

85. Boonstra TA, van Kordelaar J, Engelhart D, van Vugt JP, van der Kooij H. Asymmetries in reactive and anticipatory balance control are of similar magnitude in Parkinson's disease patients. Gait Posture 2016; 43: 108–113.

86. Smulders K, Esselink RA, De Swart BJ, Geurts AC, Bloem BR, Weerdesteyn V. Postural inflexibility in PD: does it affect compensatory stepping? Gait Posture 2014; 39: 700–706.

87. Duncan RP, Leddy AL, Cavanaugh JT, Dibble LE, Ellis TD, Ford MP, et al. Balance differences in people with Parkinson disease with and without freezing of gait. Gait Posture 2015; 42: 306–309. 11.

88. Vervoort G, Bengevoord A, Strouwen C, Bekkers EM, Heremans E, Vandenberghe W, et al. Progression of postural control and gait deficits in Parkinson's disease and freezing of gait: a longitudinal study. Parkinsonism Relat Disord 2016; 28: 73–79.

89. Earhart, G. Dance as Therapy for Individuals with Parkinson Disease. Eur J Phys Rehabil Med. 2009 June; 45(2): 231–238.

90. Hackney ME, Earhart GM. Tai Chi improves balance and mobility in people with Parkinson disease. Gait Posture 2008;28(3):456–60. [PubMed: 18378456]

91. Hackney ME, Kantorovich S, Levin R, Earhart GM. Effects of tango on functional mobility in Parkinson's disease: a preliminary study. J Neurol Phys Ther 2007;31(4):173–9. [PubMed: 18172414]

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

92. Hackney ME, Kantorovich S, Earhart GM. A study on the effects of Argentine tango as a form of partnered dance for those with Parkinson disease and the healthy elderly. Amer J Dance Ther 2007;29 (2):109–27.

93. Kwok JY, Choi KC, Chan HY. Effects of mind-body exercises on the physiological and psychosocial well-being of individuals with Parkinson's disease: A systematic review and meta-analysis. Complement Ther Med. 2016 Dec;29:121-131.

94. Westheimer O. Why dance for Parkinson's disease. Topics Ger Rehabil 2008;24(2):127–40.

95. Li F, Harmer P, Fitzgerald K, et al. Tai chi and postural stability in patients with Parkinson's disease. N Engl J Med. 2012;366: 511–519.

96. Morris ME, Iansek R, Matyas TA, et al. Ability to modulate walking cadence remains intact in Parkinson's disease. J Neurol Neurosurg Psychiatry. 1994;57:1532–1534.

97. Svehlik M, Zwick EB, Steinwender G, et al. Gait analysis in patients with Parkinson's disease off dopaminergic therapy. Arch Phys Med Rehabil. 2009;90:1880–1886.

98. Zampieri C, Salarian A, Carlson-Kuhta P, et al. The instrumented timed up and go test: potential outcome measure for disease modifying therapies in Parkinson's disease. J Neurol Neurosurg Psychiatry. 2010;81:171–176.

99. Carpinella I, Crenna P, Calabrese E, et al. Locomotor function in the early stage of Parkinson's disease. IEEE Trans Neural Syst Rehabil Eng. 2007;15:543–551.

100. Hong M, Earhart GM. Effects of medication on turning deficits in individuals with Parkinson's disease. J Neurol Phys Ther. 2010;34: 11–16.

101. Hausdorff JM, Gruendlinger L, Scollins L, et al. Deep brain stimulation effects on gait variability in Parkinson's disease. Mov Disord. 2009;24:1688–1692.

102. Schaafsma JD, Giladi N, Balash Y, et al. Gait dynamics in Parkinson's disease: relationship to Parkinsonian features, falls and response to levodopa. J Neurol Sci. 2003;212:47–53.

103. Bryant MS, Rintala DH, Hou JG, et al. Gait variability in Parkinson's disease: influence of walking speed and dopaminergic treatment. Neurol Res. 2011;33:959–964.

104. Shivitz N, Koop MM, Fahimi J, et al. Bilateral subthalamic nucleus deep brain stimulation improves certain aspects of postural control in Parkinson's disease, whereas medication does not. Mov Disord. 2006;21:1088–1097.

105. M. Tramontano, S. Bonnì, A. Martino Cinnera, F. Marchetti, C. Caltagirone, G. Koch, and A. Peppe. Blindfolded Balance Training in Patients with Parkinson's Disease: A Sensory-Motor Strategy to Improve the Gait. Parkinson's Disease. Volume 2016, Article ID 7536862, 6 pages

106. A. Peppe, C. Chiavalon, P. Pasqualetti, D. Crovato, and C. Caltagirone, "Does gait analysis quantify motor rehabilitation efficacy in Parkinson's disease patients?" Gait and Posture, vol. 26, no. 3, pp. 452–462, 2007.

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

107. Cakit BD, Saracoglu M, Genc H, Erdem HR, Inan L. The effects of incremental speed-dependent treadmill training on postural instability and fear of falling in Parkinson's disease. Clinical Rehabilitation 2007; Vol. 21, issue 8:698–705.

108. Mehrholz J, Kugler J, Storch A, Pohl M, Hirsch K, Elsner B. Treadmill training for patients with Parkinson's disease. Literature Review in Cochrane database of systematic reviews (Online) 9(9):Art. No.: CD007830. September 2015.

109. Reynard F, Terrier P. Role of visual input in the control of dynamic balance: variability and instability of gait in treadmill walking while blindfolded. Experimental Brain Research. April 2015, Volume 233, Issue 4, pp 1031–1040.

110. Jacobs JV, Nutt JG, Carlson-Kuhta P, et al. Knee trembling during freezing of gait represents multiple anticipatory postural adjustments. Exp Neurol. 2009;215:334–341.

111. Duncan RP, Leddy AL, Cavanaugh JT, Dibble LE, Ellis TD, Ford MP, Foreman KB, Earhart GM. Balance differences in people with Parkinson disease with and without freezing of gait. Gait Posture. 2015 Sep;42(3):306-9. doi: 10.1016/j.gaitpost.2015.06.007. Epub 2015 Jun 24.

112. Martina Mancini, Bastiaan R. Bloem, Fay B. Horak, Simon J.G. Lewis, Alice Nieuwboer and Jorik Nonnekes, Clinical and methodological challenges for assessing freezing of gait: Future perspectives, Movement Disorders, 34, 6, (783-790), (2019).

113. Snijders AH, Haaxma CA, Hagen YJ, Munneke M, Bloem BR.Freezer or non-freezer: clinical assessment of freezing of gait. Parkin-sonism Relat Disord 2012;18:149–154.

114. Ziegler K, Schroeteler F, Ceballos-Baumann AO, Fietzek UM. A new rating instrument to assess festination and freezing gait in par-kinsonian patients. Mov Disord 2010;25:1012–1018.21. Fietzek U, Schulz S, Ziegler K, Ceballos-Baumann AO. The FOG score detects relevant changes of gait freezing for patients and experts. Mov Disord 2018;33(Suppl 2).

115. Giladi N, Shabtai H, Simon ES, Biran S, Tal J, Korczyn AD. Construction of freezing of gait questionnaire for patients with Parkinsonism. Parkinsonism Relat Disord. 2000 Jul 1;6(3):165-170.

116. Okuma Y, Yanagisawa N. The clinical spectrum of freezing of gait in Parkinson's disease. Mov Disord. 2008;23 Suppl 2:S426-30. doi: 10.1002/mds.21934.

117. Giladi N, Shabtai H, Simon ES, Biran S, Tal J, Korczyn AD. Construction of freezing of gait questionnaire for patients with Parkinsonism. Parkinsonism Relat Disord. 2000 Jul 1;6(3):165-1170.

118. Moore O, Peretz C, Giladi N. Freezing of gait affects quality of life of peoples with Parkinson's disease beyond its relationships with mobility and gait. Mov Disord. 2007 Nov 15;22(15):2192-5.

119. Bartels AL, Balash Y, Gurevich T, Schaafsma JD, Hausdorff JM, Giladi N. Relationship between freezing of gait (FOG) and other features of Parkinson's: FOG is not correlated with bradykinesia. J Clin Neurosci. 2003 Sep;10(5):584-8.

120. Nonnekes J, Ruzicka E, Nieuwboer A, Hallett M, Fasano A,Bloem BR. Compensation strategies for gait impairments in Parkinson disease: a review. JAMA Neurol 2019 Mar 25. doi:https://doi.org/10.1001/jamaneurol.2019.0033. [

References:
One Size Fits One: Patient Centered, Impairment Based Care for Persons with Parkinson's Disease

121. Fietzek UM, Schroeteler FE, Ziegler K, Zwosta J, Ceballos-Baumann AO. Randomized cross-over trial to investigate the effi-cacy of a two-week physiotherapy programme with repetitive exer-cises of cueing to reduce the severity of freezing of gait in patientswith Parkinson's disease. Clin Rehabil 2014;28:902–911.

122. Barthel C, Nonnekes J, van Helvert M, et al. The laser shoes: a new ambulatory device to alleviate freezing of gait in Parkinson disease.Neurology 2018;90:e164–e171.

123. Nieuwboer A, Kwakkel G, Roc hester L, et al. Cueing training in thehome improves gait-related mobility in Parkinson's disease: the RES-CUE trial. J Neurol Neurosurg Psychiatry 2007; 78:134–140.

124. Haggstrom L, Darveniza P, Tisch S. Mild parkinsonian features in dystonia: Literature review, mechanisms and clinical perspectives. Parkinsonism Relat Disord. 2017 Feb;35:1-7.

125. Chaudhuri et al. International Multicenter Pilot Study of the First Comprehensive Self-Completed Nonmotor Symptoms Questionnaire for Parkinson's Disease: The NMSQuest Study. Movement Disorders. 2006: 21(7); 916–923.

126. Dirnberger G. Jahanshahi M. Executive dysfunction in Parkinson's disease: A review. 06 September 2013. https://doi.org/10.1111/jnp.12028.

127. Shirley Ryan Ability Lab. https://www.sralab.org/rehabilitation-measures/montreal-cognitive-assessment. Accessed 07/29/2019.

128. Moca Test. https://www.mocatest.org/. Accessed 07/29/19.

129. Vossius, C., Nilsen, O. B. and Larsen, J. P. (2010), Parkinson's disease and hospital admissions: frequencies, diagnoses and costs. Acta Neurologica Scandinavica, 121: 38–43. doi:10.1111/j.1600-0404.2009.01239.x

130. APTA PDEdge. http://neuropt.org/practice-resources/neurology-section-outcome-measures-recommendations/parkinson-disease. Accessed 07/29/19.

131. ParkinsonNet Guidelines. www.parkinsonnet.info/guidelines. Accessed 07/29/19.

132. Parkinson's UK Guidelines. https://www.parkinsons.org.uk/professionals/guidelines. Accessed 07/29/19.

133. Ellis, T., Cavanaugh, J.T., Earhart, G.M. Factors Associated with Exercise Behavior in People with Parkinson Disease. PHYS THER. 2011; 91:1838-1848.

134. Ellis, T. and Motl, R. Physical Activity Behavior Change in Persons with Neurological Disorders: Overview and Examples from Parkinson Disease and Multiple Sclerosis. JNPT. Volume 00, 00 2013.

135. Cress et al. Best Practices for Physical Activity Programs and Behavior Counseling in Older Adult Populations. Journal of Aging and Physical Activity, 2005; 13: 61-74.